Gray whale (*Eschrichtius robustus*) in the Mediterranean Sea: anomalous event or early sign of climate-driven distribution change?

aviad P. Scheinin 1 , dan Kerem 1 , colin D. Macleod 2 , manel gazo 3,4 , carla A. Chicote 3 and manuel castellote 5

¹Israel Marine Mammal Research and Assistance Center (IMMRAC), The Leon Recanati Institute for Maritime Studies and Department for Maritime Civilizations, The University of Haifa, Mount Carmel, Haifa 31905, Israel, ²IBES, University of Aberdeen, Aberdeen, UK, ³SUBMON—Conservation, study and awareness of the marine environment, Rabassa, 49-51. 08024 Barcelona, Spain, ⁴Department of Animal Biology, University of Barcelona, Diagonal 645, 08028 Barcelona, Spain, ⁵National Marine Mammal Laboratory, Alaska Fisheries Science Center, NOAA, Sand Point Way NE, Seattle, WA 98115 USA

On 8 May 2010, a gray whale was sighted off the Israeli Mediterranean shore and twenty-two days later, the same individual was sighted in Spanish Mediterranean waters. Since gray whales were last recorded in the North Atlantic in the 1700s, these sightings prompted much speculation about this whale's population origin. Here, we consider three hypotheses for the origin of this individual: (1) it represents a vagrant individual from the larger extant population of gray whales found in the eastern North Pacific; (2) it represents a vagrant individual from the smaller extant population found in the western North Pacific; or (3) it represents an individual from the previously thought extinct North Atlantic population. We believe that the first is the most likely, based on current population sizes, on known summer distributions, on the extent of cetacean monitoring in the North Atlantic and on the results of a performed route analysis. While it is difficult to draw conclusions from such singular events, the occurrence of this individual in the Mediterranean coincides with a shrinking of Arctic Sea ice due to climate change and suggests that climate change may allow gray whales to re-colonize the North Atlantic as ice and temperature barriers to mixing between northern North Atlantic and North Pacific biomes are reduced. Such mixing, if it were to become widespread, would have implications for many aspects of the marine conservation and ecology of these two regions.

Keywords: global warming, climate change, migration, vagrancy, sea ice, ecosystem sentinels

Submitted 29 July 2010; accepted 5 December 2010

INTRODUCTION

Gray whales (*Eschrichtius robustus*) (Lilljeborg 1861) were once found in the coastal waters of both the North Pacific and the North Atlantic. However, while historical sources indicate that gray whales were recorded on both sides of the North Atlantic until 1700, the North Atlantic population(s) seems to have been eradicated a few decades later (Lindquist, 2000). The southernmost specimen of gray whale in the eastern North Atlantic was found on the southern coast of England (Mead & Mitchell, 1984), while the southernmost western record is from Florida, USA (27°04.6′N 080°07.7′W) (Odell, 1983). While the latter record implies that a part of the North Atlantic population may have bred and calved in the lagoons of the central and south-eastern Florida coasts, the breeding and migratory habits of the gray whales that once inhabited the North Atlantic are not known.

In the North Pacific, two populations currently exist. These are a large (18,000) eastern North Pacific stock and a small (125), remnant western North Pacific stock (Swartz *et al.*, 2006). Both populations migrate northward through coastal

Corresponding author:

A.P. Scheinin Email: scheinin@research.haifa.ac.il waters in spring and summer to high latitude feeding grounds. The eastern population summers in feeding grounds in the northern Bering and southern Chukchi Seas (Moore et al., 2002), yet some whales of this population have been reported to reach longitude 174°E off the northern Chukotka coast, Russia (Berzin, 1984) and east to longitude 130°W in the Beaufort Sea, Canada (Rugh & Fraker, 1981). In autumn, this population migrates south along the eastern Pacific coast primarily to the west coast of Peninsula de Baja California, but also into the Gulf of California (Mead & Mitchell, 1984). The western population summers on feeding grounds primarily on the shallow-water shelf of north-eastern Sakhalin Island, Sea of Okhotsk (Weller et al., 2002), but also off the south-eastern coast of the Kamchatka Peninsula (Vertyankin et al., 2007). The wintering grounds of the western population are currently unknown, but may be in Asian coastal waters at latitudes as low as 20°N (Zhu, 2002).

The gray whale is the only baleen whale known to regularly consume benthic resources (Mead & Mitchell, 1984). They bottom-feed by using suction to take in food, water, and sediment, then expelling the water and sediment, while trapping the prey on the inside of their coarse baleen plates. However, gray whales are flexible foragers and respond to a variety of feeding opportunities throughout their coastal range

1

(Moore *et al.*, 2007). For example, while most gray whales forage primarily upon the shallow benthic communities of the northern seas (Nerini, 1984), whales spending the summer in lower latitudes (e.g. off Vancouver Island, Canada) can rely heavily on planktonic prey (Dunham & Duffus, 2001).

In this paper, we report on a record of an individual gray whale from two locations in the Mediterranean Sea in May 2010. Three possibilities are suggested for the population of origin of this whale:

- it represents a vagrant individual from the larger extant population of gray whales found in the eastern North Pacific;
- (2) it represents a vagrant individual from the smaller extant population found in the western North Pacific; or
- (3) it represents a surviving individual from an extant (rather than extinct) North Atlantic population.

Of the three, the first was deemed to be the most plausible, and a route analysis was done to suggest how the whale travelled to the Mediterranean and to account for its movements therein. The implications of this record for both our understanding of gray whale behaviour and the potential impacts of climate change on marine ecosystems in the North Atlantic and the North Pacific are also considered.

MATERIALS AND METHODS

Sighting records from the Mediterranean Sea

Sightings of a gray whale were recorded first off Israel and later Spain during routine cetacean surveys conducted in May 2010 in the Mediterranean Sea. During each sighting, information was gathered on the size of the individual, its body condition, behaviour such as speed and direction of travel, and photographs were taken to confirm species and individual identification, as gray whales are characterized by unique pigmentation patterns (Darling, 1984).

Investigating potential routes from the North Pacific to the North Atlantic and within the Mediterranean

In order to assess how a gray whale could have reached the Mediterranean, a geographical information system (GIS) was created in Arcview 9.3. Information entered into this database included water depth (ETOPo2, 2006), land forms and the minimum Arctic ice extent in September 2009 (from the National Snow and Ice Data Center, http://nsidc. org/data/seaice_index/), as it was assumed the odyssey began at this time. This information was then used to estimate the position and length of a number of likely migration routes from a starting point in the Bering Sea to Israel, and within the Mediterranean, for two separate scenarios. The lengths of time taken to traverse these possible routes were assessed using typical gray whale migration swimming speeds recorded in the eastern North Pacific (Mate & Harvey, 1984).

Individual identification attempts

Several eastern and western gray whale biologists were contacted and asked to review their photographic catalogues of

individual gray whales in an attempt to find a match to the gray whale sighted in the Mediterranean Sea. Western population: the Russia-US western gray whale research program catalogue, 1994 - 2009 (181 individuals), and in two catalogues of the A.V. Zhirmunsky Institute of Marine Biology, Russian Academy of Sciences (Vladivostok, Russia): (1) The Western Pacific Gray Whales of Sakhalin Island, 2002 - 2009 (177 individuals); and (2) The Pacific Gray Whales of Kamchatka Peninsula, 2004 & 2006-2009 (116 individuals). Eastern population: Cascadia Research Collective's catalogue of 'seasonal resident' gray whales that feed in the summer and autumn off California, Oregon, Washington and British Columbia (approximately 1000 individuals); the catalogue of Cape Caution in northern British Columbia (approximately 100 individuals); the catalogue of Vancouver Island (approximately 200 individuals); and the catalogue of Laguna San Ignacio Ecosystem Science Program (LSIESP) and Programa de Investigacion de Mamiferos Marinos (PRIMMA-UABCS) representing the Mexican breeding lagoons (approximately 6500 individuals).

RESULTS

Description of the sightings

Israel: a \sim 13 m long gray whale (estimated visually relative to the length of a boat) was spotted on 8 May 2010 2 km off Herzliya Marina (32°11.5′N 034°47.4′E) at 9:00 hours and followed for an hour and 10 minutes, southward along the coast. It maintained a distance of 2 km offshore, travelling at a constant speed of 5-7 km/h and performing a continuous series of short (3-5 minutes) dives occasionally displaying its flukes. A few reports of a whale near the shore had been received by the IMMRAC (Israel Marine Mammal Research and Assistance Center) stranding network two weeks before the documented sighting, suggesting that the whale was not simply transiting the area. Assessing the whale's body condition following the scale of Bradford et al. (2008), scapular condition scored 1, post-cranial condition scored 1 and lateral flank condition scored 1, for an overall inclusion in the 'poor condition' category; however, the whale did not appear to be in critical condition (i.e. near starvation) (J. Calambokidis, personal communication).

Spain: a gray whale was spotted by a sailing vessel at 16:30 hours on 30 May 2010 0.9 km away from the coast of Barcelona (41°21.32′N 002°12.2′E; north-east Spain). The whale was swimming in shallow waters of about 40 m deep, at an average speed of 7.2 km/h. The sighting lasted 20 minutes. During this period, the whale was diving constantly and displayed its flukes 4–5 times. The whale was moving southwards during the sighting period. Photographs of the tail fluke confirmed that this was the same individual previously sighted in Israel (Figure 1).

Investigating potential routes from the North Pacific to the North Atlantic

While we cannot entirely rule out the possibilities of a wintering east Pacific gray whale passing through the lock systems of the Panama Canal or around Cape Horn in order to reach the North Atlantic, nor that of a wintering west Pacific gray whale





Fig. 1. The pigmentation patterns of the flukes photographed in (A) Israel and (B) Spain, confirm that the two sightings are of the same individual. Photographs: (A) ©A. Scheinin/IMMRAC; (B) © R.Barahona/SUBMON.

traversing the South Pacific, Indian Ocean and Red Sea to enter the Mediterranean through the Suez Canal, these highly unlikely possibilities were not considered. Consequently, we considered the two most likely routes of summer-feeding whales from the Bering Sea to the North Atlantic and from there to the Mediterranean: one along the northern coast of Eurasia and the other along the northern coast of North America. Passage along either route would require that it is not blocked by sea ice. In summer 2009, of these two possible routes, the one across the northern coast of Eurasia was the most open and ice free (Figure 2), potentially making it more likely. In addition, if gray whale movements during migration are restricted to shallow shelf waters, with minimal deep-water crossings, the route via the Northwest Passage has a number of large barriers in the form of the Faroe-Iceland Rise and the Faroe-Shetland channels, while the Eurasian route does not require any deep-water crossings. On average, an eastern gray whale migrating south travels 127 km per day or 5.3 km/h (Mate & Harvey, 1984). At this speed, these two coastal routes would take a whale more than 100 days to travel between the Bering Strait and Israel. In contrast, if substantial deep-water crossings are not a barrier to migration, both routes shorten to around 85 days.

Investigating routes of movements within the Mediterranean

We considered two possible routes between the locations in Israel and Spain where this gray whale was recorded. One is a direct route over deep water, while the other is a coastal route passing only through shallow shelf waters. The former route is around 3000 km in length, while the latter is almost twice as long (Figure 2). As a result, while it would take an average swim speed of around 5.7 km/h to complete the direct route within 22 days, it would take an average swim speed of around 11 km/h to complete the coastal route. This is twice the typical migration speed for gray whales, making this route unlikely. As a result, the length of time between these two sightings suggests that this gray whale made substantial deep-water crossings during its movements.

Individual identification

The attempts to find a photographic match of the Mediterranean individual to catalogued eastern and western gray whales were unsuccessful. This might partly be due to


the fact that we have good quality images of the fluke and of the left dorsal ridge while many photo-identification catalogues are based on the right dorsal ridge and reflect surveys in shallow water areas where individuals do not regularly show their flukes.


DISCUSSION

Of the three considered points of origin, it is least likely that the gray whale observed in the Mediterranean is a remnant of a previously thought extinct North Atlantic population. In particular, given the extensive whaling and cetacean research conducted in the North Atlantic in the last 100 years, it seems unlikely that a population of shelf-foraging, large baleen whales could have survived almost 300 years without being detected.

Of the two possible North Pacific origins, the most likely origin is that of the eastern North Pacific. This population is much larger than the western Pacific population and eastern gray whale summering grounds penetrate higher into the Arctic waters to both the east and the west of the Bering Strait (Figure 2). As a result, vagrant individuals which moved further east or west than usual during the summer feeding period could end up in the North Atlantic. In contrast, the known summering grounds of the western population are at lower latitudes in the Okhotsk Sea and the eastern side of the Kamchatka Peninsula (Figure 2). This hypothesis is supported by the results of the photo-identification catalogue comparison. Given the high photographic coverage of the small western gray whale population and the more limited coverage of the larger eastern population, the fact that a match was not made suggests that the Mediterranean vagrant was not a western gray whale.

For a gray whale from the eastern population to migrate to the Mediterranean, it requires that there is an ample ice-free passage between the North Pacific and the North Atlantic, either across the top of Eurasia or North America. Therefore, the occurrence of a gray whale within the Mediterranean suggests that ice barriers to the movements of species between these two oceans have weakened to the point where passage between them is possible (MacLeod, 2010). While it is not yet clear whether such passage will become a regular event as sea ice continues to decline in response to climate change, this event suggests that such exchanges are becoming more feasible.








Fig. 2. (A) Possible routes from the typical summering ground of eastern gray whales in the Bering Sea if the whale remained in shelf waters as much as possible and restricted travel over deep water to a minimum; (B) possible routes allowing for extensive crossing of deep-water areas to minimize the distance travelled; (C) comparison of coastal versus shortest route between the sighting in Israel and the sighting near Barcelona 22 days later. Yellow lines: usual migration route for eastern Pacific gray whales; red lines: possible routes to reach the Mediterranean and to move between sighting locations; black circles: sightings locations in the Mediterranean; (1) summer feeding area for western Pacific population; (2) summer feeding area for eastern Pacific population. Background shading represents depth. The solid black is the ice extent in September 2009.

Thus, as this species and others (such as fin, humpback, and minke whales and harp, hooded, ribbon, and spotted seals) are already predisposed to disperse and become established further northward if the current overall trend toward a warmer Arctic climate continues (Moore & Huntington, 2008), the Arctic could soon become a dispersion corridor between ocean ecosystems. Therefore, the presence of a gray whale in the Mediterranean Sea concurrently with an ice-free summer Arctic passage could potentially be considered the first manifestation of such biome mixing between the North Pacific and the North Atlantic due to global climate change.

The occurrence of new species in non-native regions is thought to negatively impact marine ecosystems in many parts of the world (Ruiz et al., 1997; Hayes & Sliwa, 2003) and the spread of such species may be facilitated by changes in climate (Stachowicz et al., 2002). Specifically, the breakdown of barriers between previously isolated oceans as a result of climate change, such as that which allowed a gray whale to move into the Atlantic, could result in species moving between ocean basins with implications for the wellbeing of native marine ecosystems (Stachowicz et al., 2002). The record of this gray whale suggests that this species could potentially provide an easy-to-detect indicator of such potential exchange between the North Pacific and the North Atlantic as the Arctic ice-cap continues to shrink in response to climate change. In particular, since gray whales are a seasonally migrant species for which sea ice can act as a barrier, climate change impacts on sea ice might result in noticeable effects in their migratory behaviour, which is a conspicuous sign of ecosystem change (Laidre et al., 2008). As such, gray whales are already considered ideal sentinels of the effects of climate change on Arctic ecosystems (Moore & Huntington, 2008) and this capacity could be extended to include their use as an indicator of the ease of movement of marine mammal and other species between the Pacific and Atlantic Oceans through the currently warming Arctic corridor.

The presence of this whale within the Mediterranean also provides us with novel information about gray whales. In particular, in order to make the passage between Israel and Spain in 22 days, it is likely that this individual whale made substantial directional movements over deep-water areas where it could not forage benthically. We fully acknowledge the fact that this whale was in extraordinary circumstances, likely unfamiliar with its surroundings, such that its behaviour cannot be presumed as normal. Yet, its performance suggests that gray whales, when faced with such circumstances, can utilize and potentially forage in deep water. These observations support the concept of gray whales as the most adaptable and versatile of the mysticete species (Moore & Huntington, 2008).

Finally, the arrival of this individual to the Mediterranean also suggests that gray whales may be capable of much longer movements than are exercised during their routine migrations. If we take the breeding grounds of the eastern population as a starting point and Israel as being the southeastern extent of this individual's migration route, the whale covered a minimum distance of between $\sim\!22,000~\rm km$ and $\sim\!23,500~\rm km$, depending on the exact starting point in the eastern Pacific wintering grounds. Stevick et~al.~(2010) report on a humpback whale that travelled more than 9800 km from breeding areas in Brazil to those in Madagascar, setting a record for the largest geographical displacement ever documented in a mammal. Our results suggest

that the Mediterranean gray whale travelled more than twice that distance, resulting in what is by far the most extreme vagrancy known to have been exercised by a mammal.

ACKNOWLEDGEMENTS

The authors wish to thank Sea-Gal Sailing Club, Herzliya, and IMMRAC volunteers. Thanks are also due to Rodrigo Barahona and to the crew of ONAS sailboat for documenting the sighting off Barcelona. Special thanks are due to Phillip J. Clapham for valuable comments on the manuscript. We are grateful to all the catalogue-searching colleagues: Amanda Bradford, John Calambokidis, Dave Duffus, Alejandro Gómez-Gallardo U., Sergio Martinez, William Megill, Jorge Urbán-Ramirez, Steve Swartz, Olga Yu. Tyurneva, Yuri M. Yakovlev and Dave Weller and to two anonymous referees whose comments greatly helped to improve the manuscript.

REFERENCES

- Berzin A.A. (1984) Soviet studies on the distribution and numbers of gray whale in the Bering and Chukchi Seas from 1968 to 1982. In Jones M.L., Swartz S.L. and Leatherwood S. (eds) *The gray whale*, Eschrichtius robustus. Orlando, FL: Academic Press, pp. 409–419.
- Bradford A.L., Weller D.W., Ivashchenko Y.V., Burdin A.M. and Brownell Jr R.L. (2008) Seasonal and annual variation in body condition of western gray whales off northeastern Sakhalin Island, Russia. International Whaling Commission Scientific Committee. Paper SC/60/BRG16, 12 pp.
- Darling J.D. (1984) Gray whales off Vancouver Island, British Columbia.
 In Jones M.L. Swartz S.L. and Leatherwood S. (eds) *The gray whale*,
 Eschrichtius robustus. Orlando, FL: Academic Press, pp. 265–280.
- **Dunham J.S. and Duffus D.A.** (2001) Foraging patterns of gray whales in central Clayoquot Sound, BC, Canada. *Marine Ecology Progress Series* 223, 299 310.
- ETOPo2 (2006) Global 2' Elevations data set (U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Geophysical Data Center, version 2. 2-minute Gridded Global Relief Data. http://www.ngdc.noaa.gov/mgg/global/etopo2. html (accessed 24 May 2010).
- Hayes K.R. and Sliwa C. (2003) Identifying potential marine pests—a deductive approach applied to Australia. *Marine Pollution Bulletin* 46, 91–98.
- Laidre K.L., Stirling I., Lowry L.F., Wiig Ø., Heide-Jørgensen M.P. and Ferguson S.H. (2008) Quantifying the sensitivity of Arctic marine mammals to climate-induced habitat change. *Ecological Applications* 18 (Supplement), S97–S125.
- Lindquist O. (2000) The North Atlantic gray whale Eschrichtius robustus: an historical outline based on Icelandic, Danish–Icelandic, English and Swedish sources dating from ca 1000 AD to 1792. PhD thesis. Universities of St Andrews and Stirling, Scotland.
- **MacLeod C.D.** (2010) Global climate change, range changes and potential implications for the conservation of marine cetaceans: a review and synthesis. *Endangered Species Research* (DOI 10.3354/esr00197).
- Mate B.R. and Harvey J.T. (1984) Ocean movements of radio-tagged gray whales. In Jones M.L., Swartz S.L. and Leatherwood S. (eds) *The gray whale*, Eschrichtius robustus. Orlando, FL: Academic Press, pp. 577–589.

- Mead J.G. and Mitchell E.D. (1984) Atlantic gray whales. In Jones M.L., Swartz S.L. and Leatherwood S. (eds) *The gray whale*, Eschrichtius robustus. Orlando, FL: Academic Press, pp. 33–53.
- Moore S.E., Grebmeier J.M. and Davies J.R. (2002) Gray whale distribution relative to forage habitat in the northern Bering Sea: current conditions and retrospective summary. *Canadian Journal of Zoology* 81, 734–742.
- Moore S.E., Wynne K.M., Kinney J.C. and Grebmeier J.M. (2007) Gray whale occurrence and forage southeast of Kodiak Island, Alaska. *Marine Mammal Science* 23, 419–428.
- Moore S.E. and Huntington H.P. (2008) Arctic marine mammals and climate change: impacts and resilience. *Ecological Applications* 18, 157–165.
- Nerini M. (1984) A review of gray whale feeding ecology. In Jones M.L., Swartz S.L. and Leatherwood S. (eds) *The gray whale*, Eschrichtius robustus. Orlando, FL: Academic Press, pp. 423–450.
- Odell D.K. (1983) An Atlantic gray whale from Florida. Fifth Biennial Conference on the Biology of Marine Mammals, 27 November-1 December. Boston, MA: New England Aquarium, pp. 72-73.
- Rugh D.J. and Fraker M.A. (1981) Gray whale (Eschrichtius robustus) sightings in Eastern Beaufort Sea. Arctic 34, 186–187.
- Ruiz G.M., Carlton J.T., Grosholz E.D. and Hines A.H. (1997) Global invasions of marine and estuarine habitats by non-indigenous species: mechanisms, extent, and consequences. *American Zoologist* 37, 621–632.
- Stachowicz J.J., Terwin J.R., Whitlatch R.B. and Osman R.W. (2002)
 Linking climate change and biological invasions: ocean warming facilitates nonindigenous species invasions. *Proceedings of the National Academy of Sciences of the United States of America* 99, 15497–15500.
- Stevick P.T., Neves M.C., Johansen F., Engel M.H., Allen J., Milton C.C. Marcondes M.C.C. and Carlson C. (2010) A quarter of a world away: female humpback whale moves 10000 km between breeding areas. *Biology Letters*, doi:10.1098/rsbl.2010.0717.
- Swartz S.L., Taylor B.L. and Rugh D.J. (2006) Gray whale *Eschrichtius* robustus population and stock identity. *Mammal Review* 36, 66-84.
- Vertyankin V.V., Vladimirov V.A., Tyurneva O.Y., Yakovlev Y.M. Andreev A.V. and Burkanov V.N. (2007) Sightings of gray whales (Eschrichtius robustus) offshore eastern Kamchatka and in the northern Sea of Okhotsk, 2006. International Whaling Commission. Paper SC/59/WP6, 8 pp.
- Weller D.W., Burdin A.M., Würsig B., Taylor B.L. and Brownell Jr R.L. (2002) The western gray whale: a review of past exploitation, current status and potential threats. *Journal of Cetacean Research and Management* 4, 7–12.

and

Zhu Q. (2002) Historical records of western Pacific stock of gray whale Eschrichtius robustus in Chinese coastal waters from 1933 to 2002. International Whaling Commission Scientific Committee. Paper SC/02/WGW13, 4 pp.

Correspondence should be addressed to:

A.P. Scheinin

Israel Marine Mammal Research and Assistance Center (IMMRAC)

The Leon Recanati Institute for Maritime Studies and Department for Maritime Civilizations

The University of Haifa, Mount Carmel, Haifa 31905, Israel email: scheinin@research.haifa.ac.il